


pAd5

pAd5 is a 44 kb cosmid that contains the complete Ad5 genome between two *PacI* sites. The entire sequence of the plasmid was verified experimentally. The Ad5 genome is identical to the Adenovirus Type 5 Reference Material (GenBank [AY339865](#)), except for 3 discrepancies:

1. At position ~14080, the stretch of "A" has 14 residues in pAd5, while 13 in ARM (GenBank [AY339865](#)), 12 in Ad5 (GenBank [M73260](#)) and 17 in Ad2 (GenBank [NC_001405](#)). This homopolymer is located immediately downstream from (and includes partially) the TAA stop codon of pIIIa coding sequence.
2. At position ~29820, the stretch of "T" has 6 residues in pAd5, while 7 in ARM (GenBank [AY339865](#)), 6 in Ad5 (GenBank [M73260](#)), and 4 in Ad2 (GenBank [NC_001405](#)). In the published ARM sequence, an additional "T" would cause a frameshift mutation and premature termination of E3 10.4K (RID-alpha) protein.
3. At position ~34340, the stretch of "T" has 11 residues in pAd5, while 12 in ARM (Genbank # [AY339865](#)), 11 in Ad5 (GenBank [M73260](#)), and 14 in Ad2 (GenBank [NC_001405](#)). This homopolymer is located immediately downstream from (and includes partially) the TAA stop codon of the E4 ORF3.

The plasmid can be used for cloning purposes, as control for virus recovery by DNA transfection into helper cells, as control in Q-PCR experiments, etc...



Info Sheet	Sequence	Annotations
Product_information_pAd5.pdf (150.6 KB)	pAd5.txt (44.1 KB)	pAd5.gb (123.9 KB)